

## Peru briefing: Symptom awareness, attitudes to lung cancer and views on screening Findings from a global survey

### Overview

The Global Lung Cancer Coalition (GLCC) is an alliance of patient organisations committed to increasing awareness and understanding of lung cancer and campaigning to improve lung cancer outcomes for all.

The GLCC's Patient Charter asserts the right of all lung cancer patients to: be treated with dignity and respect; be free of blame and stigma for having the disease and to have their disease de-stigmatised; have access to quality health care; informed self-determination; physical and mental integrity; and confidentiality and privacy.

In 2023, the GLCC commissioned a multi-national study to understand attitudes towards lung cancer, levels of symptom awareness among the public and views on lung cancer screening. The study was undertaken in 29 countries: Argentina, Australia, Brazil, Bulgaria, Canada, Czech Republic, Denmark, Egypt, France, Germany, Great Britain, Greece, Italy, Israel, Japan, Mexico, Netherlands, Norway, Peru, Portugal, Republic of Ireland, Slovenia, South Africa, Spain, Sweden, Switzerland, Taiwan, Turkey and USA. Responses were received from at least 1,000 adults per country.

The study looked to build on the findings from previous studies conducted in [2010](#), [2013](#), and [2017](#), which can be found on the [GLCC website](#).

### Key findings: Peru

#### Symptom recognition

- 24.56% said they did not know any symptoms of lung cancer
- Men were more likely to report of not knowing any symptoms compared to women (27.43% v 21.58%)
- People aged 18-24 were most likely to report not knowing any symptoms (32.52%)
- People who have never smoked were more likely to report not knowing any symptoms compared to people who currently smoke/have smoked (29.60% v 22.24%)

#### Attitudes towards lung cancer

- 43.88% of people agreed with the statement 'I have less sympathy for people with lung cancer than other types of cancer'; this compares to 42.62% globally
- Men had less sympathy for people with lung cancer than women (50.10% v 37.43%)
- People aged 25-34 had the least sympathy for people with lung cancer than other types of cancer, while individuals aged 45-54 had the most (49.61% v 32.48%)

### **The GLCC is calling on governments and health systems across the world to:**

- Increase recognition of lung cancer symptoms through targeted public awareness campaigns
- Improve public education and understanding of lung cancer and its causes to reduce stigma
- Highlight the importance of early diagnosis and treatment in improving survival to encourage more people to seek help early
- Implement national lung cancer screening programme to help people get diagnosed and treatment as early as possible

- People who currently smoke/have smoked were less sympathetic than non-smokers (44.76% v 42.37%)

### Views on lung cancer screening

- 93.20% people agreed there should be a national lung cancer screening programme to help people with lung cancer get diagnosed early
- There was near-equal support for lung cancer screening programmes between men and women (93.14% v 93.27%)
- Support was highest among respondents aged 45-54, and lowest among respondents aged 18-24 (97.44% v 88.83%)
- People who currently smoke/have smoked were marginally more supportive of lung cancer screening programmes than people who have never smoked (93.83% v 92.52%)

### Methodology

The GLCC commissioned the market research agency Censuswide to undertake an online survey of adults across 29 countries between 3 July 2023 – 6 July 2023. The survey samples included at least 1,000 adults (per country) and data was weighted using age, gender and region to produce a representative sample in each country. Respondents were asked the following questions:

- Q1. There are many warning signs and symptoms of lung cancer. Please list as many symptoms of lung cancer as you can think of [a tick box was provided for 'Don't know']
- Q2. Lung cancer is mainly caused by smoking cigarettes and other tobacco products. To what extent do you agree or disagree with the following statement: I have less sympathy for people with lung cancer than for other types of cancer
- Q3. To what extent do you agree or disagree with the following statement: There should be a national lung cancer screening programme to help people with lung cancer get diagnosed early

### Further analysis

#### Symptom recognition

Raising awareness of lung cancer symptoms is vital because treatment is more likely to be successful if lung cancer is diagnosed early.

The study found that symptom awareness is generally poor. Within Peru, 24.56% of respondents reported not knowing any symptoms, showing that while marked progress has been made, more needs to be done to support greater awareness of lung cancer symptoms.

Chart 1. Percentage of respondents who indicated they did not know any symptoms by country




Chart 2. Percentage of respondents in Peru who indicated they did not know any symptoms by age





Chart 3. Percentage of respondents in Peru who indicated they did not know any symptoms by smoking status



### Attitudes towards lung cancer

More than eight out of ten cases of lung cancer are linked to smoking creating stigma around the disease. This is a problem not only because thousands of people develop lung cancer who have never smoked, but also because people who have smoked may feel guilty and delay talking their doctor about potential symptoms.

The GLCC believes that everyone - no matter what the cause of their cancer - deserves to have high quality treatment and care. No one deserves to have lung cancer. If lung cancer can be picked up early then it improves people's chance of successful treatment and survival.

To understand attitudes to lung cancer, participants were told that: “Lung cancer is mainly caused by smoking cigarettes and other tobacco products”. They were then asked whether or not they have less sympathy for people with lung cancer than other types of cancer.

In Peru, 43.88% agreed with the statement that they had less sympathy for people with lung cancer.

Chart 4. Percentage of respondents who agree with the statement they have less sympathy for people with lung cancer than for other types of cancer by country




Chart 5. Percentage of respondents in Peru who agree with the statement they have less sympathy for people with lung cancer than for other types of cancer by age





Chart 6. Percentage of respondents in the Peru who agree with the statement they have less sympathy for people with lung cancer than for other types of cancer by smoking status



## Views on lung cancer screening


A national lung cancer screening programme is a key measure to reduce the huge burden that the disease currently imposes on the individuals affected, their families and the country and healthcare system as a whole.

Lung cancer screening with a non-invasive scan of the lungs (low-dose computed tomography [LDCT]) saves lives by detecting the disease at an early stage when it is potentially curable. A large clinical trial of LDCT screening showed a 24% reduction in lung cancer mortality in men and a 33% reduction in women at 10 years of follow-up compared to no screening (de Koning 2020).


It is encouraging to see that, in Peru, there is widespread support for a national lung cancer screening programme, with 93.20% of participants responding in favour.

At present only a few countries have national lung cancer screening programmes. Given the impact they can have and the support for them globally, the GLCC calls on policymakers around the world to implement national lung cancer screening programmes to help people with lung cancer get diagnosed as early as possible.


*Chart 7. Percentage of respondents who agree there should be a national lung cancer screening programme to help people with lung cancer get diagnosed early by country*



*Chart 8. Percentage of respondents in the Peru who agree there should be a national lung cancer screening programme to help people with lung cancer get diagnosed early by age*



*Chart 9. Percentage of respondents in Peru who agree there should be a national lung cancer screening programme to help people with lung cancer get diagnosed early by smoking status*



## More information

Established in 2001, the GLCC comprises 43 non-government patient organisations from Argentina, Australia, Brazil, Bulgaria, Canada, Czech Republic, Denmark, Egypt, France, Germany, Greece, Ireland, Israel, Italy, Japan, Mexico, Netherlands, Norway, Peru, Portugal, Russia, Slovenia, South Africa, Spain, Sweden, Switzerland, Taiwan, Turkey, UK and US.

The GLCC promotes global understanding of lung cancer and the right of patients to effective early detection, better treatment and supportive care. By serving as the international voice of lung cancer patients, the GLCC is committed to improving disease outcomes for all.

For more information about this study and the work of the GLCC please visit: [www.lungcancercoalition.org](http://www.lungcancercoalition.org) or email our secretariat at: [glcc@roycastle.org](mailto:glcc@roycastle.org)